

GROPRO

STOMP SLUG

LIQUID CONCENTRATE

ACTIVE INGREDIENT:

Garlic Oil (8000-78-0)..... 2.00%

OTHER INGREDIENTS:* 98.00%**TOTAL:** 100.00%

*Isopropyl Alcohol, Tea Saponin, Water

MANUFACTURED BY:**GROPRO**12600 West Frontage Road,
Burnsville MN 55337

Phone: 1-833-476-7761

www.groproag.com

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Precautionary Statements

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle (If you do not understand the label, find someone to explain it to you in detail)

How can we help? 1-833-476-7761

ENVIRONMENTALLY SAFE
When Used as Directed

NET CONTENTS:

LOT#: _____

■ 2.5 Gal (9.5 L) ■ 30 Gal (113.6 L) ■ 264 gal (999.3 L)

FIRST AID

IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything to an unconscious person.

IF INHALED: • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.

POISONING HOTLINE

Seek medical care as soon as possible after exposure. For incidents involving animals, seek veterinary care as soon as possible after exposure. Have the product container or label with you when calling a poison control center or doctor or veterinarian, or when going for treatment. For information on this pesticide product (including health concerns, medical emergencies, treatment advice, or pesticide incidents), call National Pesticide Information Center at 1-800-858-7378.

Tea saccharides is a natural material, which is safe for humans.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Protective eye wear
- Chemical resistant gloves
- Chemical resistant apron (for cleaning, mixing, or loading)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. This pesticide is non-toxic to birds and mammals. Liquid on soil surface is not hazardous to terrestrial wildlife.

STORAGE AND DISPOSAL

STORAGE AND DISPOSAL Prohibitions: Do not contaminate water, food or feed by storage and disposal. **Pesticide Storage:** Store this product in original container in a cool, secure location out of reach of children and pets. Protect from freezing. **Pesticide Disposal:** To avoid wastes, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often

such programs are run by state or local governments or by industry).

Container Disposal: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Triple rinse container (or equivalent) promptly after emptying. Offer for recycling, if available or offer for reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or incineration.

-For over 2.5, 30, 264 gallons containers; triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. This product is intended solely for use on agricultural crops grown for commercial or research purposes. This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

REENTRY: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 0 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated areas without restriction if there will be no contact with anything that has been treated.

PRECAUTIONS AND RESTRICTIONS

- This product is intended solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication operations. This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.
- Broadcast applications and applications of this product directed to plant parts are permitted.
- Do not apply to dry soil. Thoroughly water areas to be treated before application.
- This product may be tank mixed with other compatible chemicals or fertilizers. •Evening applications are preferred.

MIXING: Shake Stomp SLUG CONTROL well before mixing. Thoroughly mix spray solution before application. Use entire mixed solution at one time. Do not store mixed solution.

ARTICHOKE - perennial: Thoroughly water ground areas to be treated before application. Do not apply in such a manner as to cause drift of product. For best results, apply in the evening, and in enough water for thorough coverage on foliage.

SLUGS AND SNAILS: Apply 64-96 fl. oz. per treated acre in sufficient water for thorough coverage.

ARTICHOKE – annual: Thoroughly water ground areas to be treated before application. Do not apply in such a manner as to cause drift of product. For best results, apply in the evening, and in enough water for thorough coverage on foliage.

SLUGS AND SNAILS: Apply 64-96 fl. oz. per treated acre in sufficient water for thorough coverage.

BERRIES: Caneberries (bingleberry, black raspberry, blackberry, boysenberry, dewberry, lowberry, marionberry, olallieberry, red raspberry, youngberry) and Other Berries (cranberry, currant, elderberry, gooseberry, huckleberry, loganberry, lingonberry, juneberry, salal), Blueberries. Thoroughly water areas to be treated before application. Thoroughly water ground areas to be treated before application. Do not apply in such a manner as to cause drift of product. For best results, apply in the evening, and in enough water for thorough coverage on foliage.

SLUGS AND SNAILS: Apply 64-96 fl. oz. per treated acre in sufficient water for thorough.

CITRUS: Make application when foliage and fruit are completely dry but soil is wet following an irrigation. Apply to soil under trees. Use equipment that applies directly to soil surface. Thoroughly water ground areas to be treated before application. Do not apply in such a manner as to cause drift of product. For best results, apply in the evening, and in enough water for thorough coverage on foliage.

SLUGS AND SNAILS: Apply 64-96 fl. oz. per treated acre in sufficient water for thorough coverage.

Lettuce, Cole Crops and Other Leafy Greens (broccoli, Brussels sprouts, cabbage, cauliflower, cavalo, celery, broccolo, collards, kale, kohlrabi, mizuna, mustard greens, spinach, swiss chard, rape greens) **Tomatoes, Rhubarb and Strawberries.** Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to the soil surface and plant foliage. Thoroughly water ground areas to be treated before application. Do not apply in such a manner as to cause drift of product. For best results, apply in the evening, and in enough water for thorough coverage on foliage.

For best results, apply in the evening. **SLUGS AND SNAILS:** Apply 64-96 fl. oz. per treated acre in sufficient water for thorough coverage.

ORNAMENTALS: Thoroughly water areas to be treated before application. Use equipment that applies directly to the soil surface. Apply evenly over infested areas and along flower beds adjacent to border or ground cover plants. Do not apply in such a manner as to cause drift of product onto plants. NOTE: Some slight tip burning has been noticed on extremely tender orchids. For best results, apply in the evening.

SLUGS AND SNAILS: Apply 64-96 fl. oz. per treated acre in sufficient water for thorough coverage OR 2.25 fl. oz. in ten gallons of water per treated 1,000 sq. ft. OR 3/4 tablespoon in two gallons of water per 200 sq. ft.

PRICKLY PEAR CACTUS: The ground onto which the Stomp Slug Control is applied must be wet (via rainfall or irrigation). The crop may be harvested on the day of application.

SLUGS AND SNAILS: Apply Stomp Slug Control to the ground in a line on each side of the crop, 1-3 feet from the base of the plants, at a rate of 64-96 fluid ounces per treated acre in sufficient water for thorough coverage.

GRASS GROWN FOR SEED: Uniformly spray directly on grass grown for seed. Withhold heavy watering for at least 2 days after treatment. Keep children and pets off the treated area until spray has dried.

SLUGS AND SNAILS: Apply 64-96 fl. oz. per treated acre in sufficient water for thorough coverage OR 1.5 fl. oz. in ten gallons of water per treated 1,000 sq. ft.

WATERCRESS: •Application rate: 118 fl. oz. per acre.

- Do not make more than four applications per harvested crop.
- The first application can be made following planting a new crop or following the harvest and start of a return crop. The second application can be made up to harvest.
- A maximum of 8 applications per year can be made to a field.

PRE-HARVEST INTERVAL: 0 days.

USE DIRECTIONS

- Fields to be treated with Stomp Slug Control must be drained for a minimum period of 24 hours prior to treatment and must remain drained for 24 hours following the treatment.
- Apply to the crop canopy.
- For optimum efficiency, apply Stomp Slug Control in the morning while dew is still on the plants and snails are visible on the plants.
- Watering restrictions: avoid irrigation or addition of water until Stomp Slug Control is dry.
- The average depth of water in the watercress beds should be between 1 to 1.5 inches while watercress is in production.
- Make applications by ground sprayer equipment with a minimum of 50 gallons of water per acre. Applications can also be made through chemigation system. When chemigation is used, follow the directions for chemigation system.

CHEMIGATION USE INSTRUCTIONS

Apply Stomp Slug Control only through solid set, micro sprinkler, and drip irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, in the crop can result from non-uniform distribution of material. If you have questions about calibration, you should contact State Extension Service specialist. Do not connect an irrigation system used for pesticide application to a public water system.

OPERATING INSTRUCTIONS:

- 1.** The irrigation system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- 2.** The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
- 3.** The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4.** The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5.** System must use a metering pump effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 6.** Do not apply when wind speed favors drift beyond the area to be treated.
- 7.** If using solid sets or sprinklers part circle sprinklers should be installed at the periphery of the fields to prevent an off-target application.
- 8.** Stomp Slug Control must be applied on the schedule specified in the specific crop use directions, and not according to the irrigation schedule.
- 9.** Do not inject Stomp Slug Control at full strength. Dilute in adequate amount of water prior to injecting in the irrigation system for chemigation.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions of Warranties and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer or Warranties and Limitation of Liability.

Conditions: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness of other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GROPRO Inc. All such risks shall be assumed by the user or buyer.

Disclaimer or Warranties: To the extent consistent with applicable law, GROPRO Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of GROPRO Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, GROPRO Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

Limitations of Liability: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at GROPRO Inc's election, the replacement of product.